

GENERALIZED MODELS AND LOCAL INVARIANTS OF KOHN-NIRENBERG DOMAINS

MARTIN KOLÁŘ

ABSTRACT. The main obstruction for constructing holomorphic reproducing kernels of Cauchy type on weakly pseudoconvex domains is the Kohn-Nirenberg phenomenon, i.e., nonexistence of supporting functions and local nonconvexifiability. In this paper we give an explicit verifiable characterization of weakly pseudoconvex but locally nonconvexifiable hypersurfaces of finite type in dimension two. It is expressed in terms of a generalized model, which captures local geometry of the hypersurface both in the complex tangential and nontangential directions. As an application we obtain a new class of nonconvexifiable pseudoconvex hypersurfaces with convex models.

1. INTRODUCTION

Although local convexity of the boundary of a weakly pseudoconvex domain is not a biholomorphic invariant, it is often used as an assumption which provides useful tools to study biholomorphically invariant objects (see e.g. [4], [6], [13]). The natural invariant condition for results of this kind is then just that the domain be locally biholomorphic to a convex domain. We consider here the problem of finding an explicit characterization of such domains.

Let $D \subseteq \mathbb{C}^n$ be a domain with smooth boundary M and p be a point on M . Let ψ be a local defining function in a neighbourhood U of p . Recall that D is pseudoconvex in U if for all $q \in M \cap U$ the Levi form

$$\mathcal{L}_q(\zeta) = \sum_{i,j=1}^n \frac{\partial^2 \psi}{\partial z_i \partial \bar{z}_j}(q) \zeta_i \bar{\zeta}_j$$

is nonnegative on the complex tangent space to M at q , i.e. on complex vectors $\zeta = (\zeta_1, \zeta_2, \dots, \zeta_n)$ satisfying $\sum_{i=1}^n \frac{\partial \psi}{\partial z_i}(q) \zeta_i = 0$. When D is strictly pseudoconvex (the Levi form is positive definite at each point), fundamental results of Henkin and Ramirez provide a holomorphic reproducing kernel, a direct analog of Cauchy's integral kernel from one variable. The construction is based on holomorphic supporting functions $h_q(z)$, defined in a neighbourhood of each $q \in M$, whose zero set intersects $\bar{D}$ only at q . The reciprocal of a supporting function is the analog of the fundamental function

The author was supported by a grant of the GA ĆR no. 201/05/2117.

$\frac{1}{z-q}$ from one complex variable. The existence of such functions is a consequence of the following stronger result, known as Narasimhan's lemma.

Theorem 1.1 (Narasimhan). *Let $D \subseteq \mathbb{C}^n$ be a smoothly bounded domain, which is strictly pseudoconvex in a neighbourhood of a boundary point p . Then there exist local holomorphic coordinates around p in which ∂D is strictly convex.*

In 1973, Kohn and Nirenberg found an example which shows that local convexifiability does not in general extend to weakly pseudoconvex domains. It disproved the popular conjecture that pseudoconvexity is equivalent to local convexifiability. Their example is the domain

$$D_0 = \{(z, w) \in \mathbb{C}^2 \mid \operatorname{Im} w > |z|^8 + \frac{15}{7}|z|^2 \operatorname{Re} z^6\}$$

which does not admit a holomorphic supporting function at the origin, and therefore is not convexifiable. As a consequence, in connection with the efforts to extend Henkin's construction to (at least some) weakly pseudoconvex domains, it became an important problem to characterize domains for which this phenomenon occurs.

Local convexifiability of finite type domains turns out to be much stronger than mere existence of supporting functions in all neighbouring points. The problem of explicit characterization of locally convexifiable domains in complex dimension two was considered in [10] and [11], with a satisfactory answer on the level of model domains. The results are based on exact conditions computed for analogs of the Kohn-Nirenberg example (see Proposition C below). They do not take into account behavior of the defining function in the complex non-tangential direction. Moreover, convexifiability of the model domain is necessary, but not sufficient for convexifiability of the domain itself. Hence the possibility of other obstructions to convexifiability remained open.

Our aim in this paper is to complete the results of [10], [11], by introducing a generalized model domain which (up to a borderline case) carries the information about local convexifiability of the domain itself. In this way, the problem which apriori requires to consider the whole infinite dimensional group of local biholomorphisms is reduced to a simple finite dimensional problem, which is often solved by an application of the explicit results of [10].

As an application, we find a new class of nonconvexifiable domains. While all previous examples are on the level of model domains, we describe nonconvexifiable pseudoconvex domains whose model domains at each point are convex.

In Section 2 we use weighted coordinates to introduce the generalized model domain. The construction is similar to Catlin's definition of multitype in $\mathbb{C}^n$ (see [1]), where only complex tangential variables are considered. The weight of the complex nontangential direction provides a local biholomorphic invariant with rational values. Section 3 gives two complementary conditions characterizing local convexifiability. The proof of the sufficient condition relies on weighted homogeneity of the generalized model. The main tool in proving the necessary condition is the technique used in [3] for analyzing biholomorphic transformations in weighted coordinates. Section 4 reviews the explicit conditions for local convexifiability derived in [10], [11]. In Section 5 we describe a new class of examples of nonconvexifiable domains which have convex models at each point.

A part of this work was done while the author was visiting the ESI program Complex Analysis, Operator Theory, and Applications to Mathematical Physics. He would like to thank the organizers, Friedrich Haslinger, Emil Straube and Hans Upmeyer for their invitation and hospitality, and for the support recieved from ESI.

2. GENERALIZED MODEL DOMAINS

We will consider a pseudoconvex domain $D \subseteq \mathbb{C}^2$ with smooth boundary M , and a point $p \in M$ of finite type k , where $k > 2$. Recall that the type of a point measures the maximal order of contact of M with complex curves passing through p . Strongly pseudoconvex points are points of type two. It follows from pseudoconvexity that k is an even integer.

We will use local holomorphic coordinates (z, w) , $z = x + iy$, $w = u + iv$, centered at p , such that the positive v -axis gives the inner normal direction to D at p and write the defining equation for M as

$$v = F(z, \bar{z}, u).$$

It follows directly from the definition of finite type (see [7]) that we can choose the coordinates so that the equation takes form

$$(2.1) \quad v = P_1(z, \bar{z}) + o(|z|^k, u),$$

where P_1 is a real valued homogeneous polynomial of degree k

$$(2.2) \quad P_1(z, \bar{z}) = a_0|z|^k + \sum_{j=2, \dots, k} |z|^{k-j} \operatorname{Re}(a_j z^j)$$

for some $a_j \in \mathbb{C}$ and $a_0 \in \mathbb{R}^+$. The hypersurface

$$(2.3) \quad M_D = \{(z, w) \in \mathbb{C}^2 \mid v = P_1(z, \bar{z})\}$$

is called a model hypersurface to M at p . It is determined uniquely up to a linear change of variables and addition of a harmonic term $\operatorname{Re} \alpha z^k$ (see e.g. [11]).

Definition 2.1. M is called locally convexifiable at p if there exist local holomorphic coordinates in a full neighbourhood of p such that M is convex with respect to the underlying linear space.

The polynomial P_1 in (2.1) captures local behavior of M in the complex tangential direction. In order to study convexifiability of M near p we wish to take into account also behavior in complex nontangential directions, involving the variable u . A natural tool for this is provided by weighted coordinates. We will assign weights to the coordinates z , w and u .

Definition 2.2. A weight vector is a pair of rational numbers $\lambda = (\lambda_1, \lambda_2)$ such that $\lambda_1 = \frac{1}{n}$ for some $n \in \mathbb{N}$ and there exist integers k_1, k_2 , with $k_2 > 0$ such that $k_1 \lambda_1 + k_2 \lambda_2 = 1$.

The weight of monomials $c_{ijk} z^i \bar{z}^j u^k$ and $d_{ij} z^i w^j$ is defined to be $(i+j)\lambda_1 + k\lambda_2$ and $i\lambda_1 + k\lambda_2$, respectively. A real valued polynomial $Q(z, \bar{z}, u)$ is λ -homogeneous of weight γ if it is a sum of monomials of weight γ , and similarly for a holomorphic polynomial $P(z, w)$. The weight of a function $Q(z, w)$ is the lowest of the weights of the nonzero terms in its Taylor expansion at the origin (when working in weighted coordinates it

is helpful to use the two dimensional lattice $\mathbb{Z}^+ \oplus \mathbb{Z}^+$, where a monomial $z^a \bar{z}^b u^c$ in the Taylor expansion of F is assigned the lattice point $(a + b, c)$.

Given the point $p \in M$, we first assign weight to the variable z , equal to the reciprocal of the type of the point, $\lambda_1 = \frac{1}{k}$. The weight of w and u will be denoted by μ , and is determined as follows. We will call a weight vector $\lambda = (\frac{1}{k}, \lambda_2)$ allowable, if there exist local holomorphic coordinates such that the defining equation can be written in the form

$$v = P(z, \bar{z}, u) + o_{wt}(1),$$

where

$$P(z, \bar{z}, u) = \sum_{\frac{m}{k} + \lambda_2 l = 1} \sum_{j=0}^m a_{mjl} z^j \bar{z}^{m-j} u^l$$

is a λ -homogeneous polynomial of weight one, and $o_{wt}(1)$ denotes terms of weight bigger than one with respect to the weight vector $(\frac{1}{k}, \lambda_2)$.

Clearly, for any real $\delta > 0$ there are only finitely many rational numbers λ_2 bigger than δ such that $(\frac{1}{k}, \lambda_2)$ is a weight. When the set of allowable values for λ_2 is finite, we set μ to be the smallest of them. If the set is not bounded away from zero, we set $\mu = 0$. Using the change of variables formula (3.3) below, it is straightforward to determine the value of μ .

When $\mu > 0$, we fix local holomorphic coordinates (z, w) which correspond to the weight $\lambda = (\frac{1}{k}, \mu)$. Hence the defining equation has form

$$(2.4) \quad v = \sum_{\frac{m}{k} + \mu l = 1} \sum_{j=0}^m a_{mjl} z^j \bar{z}^{m-j} u^l + o_{wt}(1),$$

where, as before, the leading term will be denoted by $P(z, \bar{z}, u)$. The hypersurface given by

$$v = P(z, \bar{z}, u)$$

will be called a generalized model to M at p . If $\mu = 0$, a generalized model is defined to be

$$v = P_1(z, \bar{z}),$$

hence in this case it coincides with the standard model hypersurface.

3. CHARACTERIZATIONS OF CONVEXIFIABILITY

Let $\mathbb{H}_{k,\mu}$ denote the set of all real valued λ -homogeneous polynomials in z, u of weight one which are harmonic in the z variable. Clearly, $h(z, u) \in \mathbb{H}_{k,\mu}$ if and only if

$$h(z, u) = \operatorname{Re} \left(\sum_{\frac{m}{k} + \mu l = 1} \alpha_m z^m u^l \right)$$

for some $\alpha_m \in \mathbb{C}$.

For $(z, u) \in \mathbb{C} \times \mathbb{R}$ and $\zeta \in \mathbb{C}$ we will use the following notation for the value of the real Hessian restricted to the z -direction:

$$D_z^2 P(z, u; \zeta) = \sum_{i,j=1}^2 \frac{\partial^2 P}{\partial x_i \partial x_j}(z, u) \xi_i \xi_j,$$

where $z = x_1 + ix_2$ and $\zeta = \xi_1 + i\xi_2$. Since P is weighted homogeneous, positivity of $D_z^2 P$ is determined by its restriction to the set $S_2 \times S_1$, where $S_2 = \{(z, u) \in \mathbb{C} \times \mathbb{R} : |z|^2 + u^2 = 1\}$ and $S_1 = \{\zeta \in \mathbb{C} : |\zeta| = 1\}$.

Our aim is to prove the following pair of conditions for convexifiability of M .

Proposition A. *If there exists $h \in \mathbb{H}_{k,\mu}$ such that $D_z^2(P + h) > 0$ on $S_2 \times S_1$, then M is locally convexifiable.*

Proof. Let $h \in \mathbb{H}_{k,\mu}$ be such a function and let $\tilde{P} = P + h$. Considering first the condition on the u -axis, it follows from the assumption that $\frac{1}{\mu}(1 - \frac{2}{k})$ is an even integer and that P contains a term $|z|^2 u^{\frac{1}{\mu}(1 - \frac{2}{k})}$ with a positive coefficient. By homogeneity, we have

$$(3.1) \quad D_z^2 \tilde{P}(z, u; \zeta) \geq \epsilon(|z|^{k-2} + |u|^{\frac{1}{\mu}(1 - \frac{2}{k})})|\zeta|^2$$

for a sufficiently small $\epsilon > 0$. Consider the change of coordinates

$$w^* = w + iw^2, \quad z^* = z,$$

and denote $v^* = F^*(z^*, \bar{z}^*, u^*)$ the defining equation in the new coordinates. We get

$$F^*(z, \bar{z}, u - 2uF(z, \bar{z}, u)) = F(z, \bar{z}, u) + u^2 - (F(z, \bar{z}, u))^2.$$

Comparing terms of weight ≤ 1 on both sides we obtain

$$F^*(z^*, \bar{z}^*, u^*) = \tilde{P}(z^*, \bar{z}^*, u^*) + (u^*)^2 + o_{wt}(1).$$

We will show that F^* is a convex function in a neighbourhood of the origin. Denote $P^*(z, \bar{z}, u) = \tilde{P}(z, \bar{z}, u) + u^2$ (dropping stars on variables) and consider its 3×3 Hessian matrix with respect to the variables x, y, u . The 2×2 submatrix of the Hessian formed by the derivatives with respect to x and y satisfies (3.1), hence is positive semidefinite. It remains to prove that the determinant of the Hessian is also nonnegative in a neighbourhood of the origin. We calculate the weights of the terms entering into the determinant. On the one hand, $wt(P_{uu}^* P_{xx}^* P_{yy}^*) = wt(P_{uu}^* P_{xy}^* P_{yx}^*) = 2 - \frac{4}{k}$. On the other hand, $wt((P_{xu}^*)^2 P_{yy}^*) = 3 - 2\mu - \frac{4}{k}$ and the same for all the remaining terms. Since $k > 2$ and $\mu < \frac{1}{2}$, we have $3 - 2\mu - \frac{4}{k} > 2 - \frac{4}{k}$. By (3.1), the determinant is nonnegative in a neighbourhood of the origin. $\square$

Proposition B. *If M is locally convexifiable, then there exists $h \in \mathbb{H}_{k,\mu}$ such that $D_z^2(P + h) \geq 0$ on $S_2 \times S_1$.*

Proof. We will assume that $D_z^2(P + h)$ is not nonnegative on $S_2 \times S_1$ for any $h \in \mathbb{H}_{k,\mu}$, and show that M cannot be locally convex in any other coordinates. If $\mu = 0$, the

claim follows from Lemma 3 in [11], so we assume $\mu > 0$. Consider a holomorphic transformation

$$(3.2) \quad \begin{aligned} z^* &= z + g(z, w) \\ w^* &= w + f(z, w). \end{aligned}$$

We may restrict ourselves to transformations which preserves the above description of M , so $v^* = 0$ is tangent to M at the origin and the positive v^* -axis points inside. Let F^* be the function describing M in new coordinates. Substituting (3.2) into $v^* = F^*(z^*, \bar{z}^*, u^*)$, we get the change of variables formula relating F^* and F , f , g ,

$$(3.3) \quad \begin{aligned} F^*(z + g(z, u + iF(z, \bar{z}, u)), \overline{z + g(z, u + iF(z, \bar{z}, u))}, u + \operatorname{Re} f(z, u + iF(z, \bar{z}, u))) = \\ = F(z, \bar{z}, u) + \operatorname{Im} f(z, u + \operatorname{Re} f(z, u + iF(z, \bar{z}, u))). \end{aligned}$$

Without any loss of generality we assume that the linear part of the transformation is partly normalized and together require that

$$f = 0, \quad g = 0, \quad f_z = 0, \quad g_z = 0 \quad f_w = 0 \quad \text{at} \quad z = w = 0.$$

Germes of transformations which satisfy this condition form a group, which will be denoted by $\mathcal{G}$.

We will call $g(z, w)$ in (3.2) strictly superhomogeneous if $wt(g) > wt(z) = \frac{1}{k}$. By $\mathcal{G}_1$ we will denote the subgroup of $\mathcal{G}$ formed by germes of transformations (3.2) for which g is strictly superhomogeneous.

Next we show that every $G \in \mathcal{G}$ can be factored uniquely as $G = \tau \circ S$, where $\tau \in \mathcal{G}_1$ and S is a polynomial transformation of the form

$$(3.4) \quad \begin{aligned} z^{**} &= z^* + \sum_{j=1}^{\lfloor \frac{1}{k\mu} \rfloor} \delta_j (w^*)^j, \\ w^{**} &= w^*. \end{aligned}$$

Indeed, write $G \in \mathcal{G}$ as

$$\begin{aligned} z^* &= z + \sum_{j=1}^{\infty} \alpha_j w^j + \sum_{i=1, j=0}^{\infty} \beta_{ij} z^i w^j, \\ w^* &= w + \sum_{j=2}^{\infty} \epsilon_j w^j + \sum_{i=1, j=0}^{\infty} \gamma_{ij} z^i w^j \end{aligned}$$

and compose it with a transformation S of the form (3.4). We claim there exist uniquely determined numbers δ_j such that $G \circ S \in \mathcal{G}_1$. In order to see this, denote the composition by

$$z^{**} = z + g^{**}(z, w), \quad w^{**} = w + f^{**}(z, w),$$

where

$$g^{**}(z, w) = \sum_{j=1}^{\infty} \alpha_j w^j + \sum_{i=1, j=0}^{\infty} \beta_{ij} z^i w^j + \sum_{m=1}^{\lfloor \frac{1}{k\mu} \rfloor} \delta_m (w + \sum_{j=2}^{\infty} \epsilon_j w^j + \sum_{i, j=0}^{\infty} \gamma_{ij} z^i w^j)^m$$

$$f^{**}(z, w) = \sum_{j=1}^{\infty} \epsilon_j w^j + \sum_{i=1, j=0}^{\infty} \gamma_{ij} z^i w^j.$$

There is a unique way to choose the coefficients δ_j so that g^{**} is strictly superhomogeneous, namely $\delta_1 = -\alpha_1$, $\delta_2 = -\alpha_2 - \delta_1 \epsilon_2$, and so on up to index $\lfloor \frac{1}{k\mu} \rfloor$. Since the inverse of S is of the same form as S , we obtain the claimed decomposition.

Now consider the transformation by a representant of a fixed element $G_0 \in \mathcal{G}$ and use the above decomposition, $G_0 = \tau_0 \circ S_0$. Let τ_0 be of the form (3.2) and consider (3.3) for terms of weight less or equal to one.

Now we consider two cases. When f contains only terms of weight greater or equal to one, then τ_0 preserves form (2.4), merely adding harmonic terms to P . So it's enough to consider the effect of S_0 , given by

$$(3.5) \quad \begin{aligned} z^* &= z + \sum_{j=1}^{\lfloor \frac{1}{k\mu} \rfloor} \delta_j w^j, \\ w^* &= w. \end{aligned}$$

By assumption, there exists a smooth curve γ through the origin in $\mathbb{C} \times \mathbb{R}$ and a vector $\zeta_0 \neq 0$ such that in a neighbourhood of p

$$(3.6) \quad D_z^2 P(z, u; \zeta_0) \leq -\epsilon(|z|^{k-2} + |u|^{\frac{1}{\mu}(1-\frac{2}{k})})|\zeta_0|^2$$

for $(z, u) \in \gamma$ and a sufficiently small $\epsilon > 0$. Consider terms of degree less or equal to $s+2$ in F , where $s = (1 - \frac{2}{k})\frac{1}{\mu}$. They will be denoted by $T(z, \bar{z}, u)$, so $F = T(z, \bar{z}, u) + o(s+2)$. Here the ordinary (nonweighted) degree is used, $o(s+2)$ denoting terms of degree greater than $s+2$. T contains all terms which after applying S_0 can influence terms of weight less or equal to one in F^* . The estimate (3.6) holds also for $D_z^2 T$, perhaps with a smaller ϵ and a smaller neighbourhood of p . After applying S_0 (and dropping stars on variables) we consider terms of weight less or equal to one in F^* , and denote them by Q^* . By (3.3) we obtain, modulo terms $o_{wt}(1)$,

$$Q^*(z, \bar{z}, u) = T(z + \sum_{m=1}^{\lfloor \frac{1}{k\mu} \rfloor} \delta_m u^m, \bar{z} + \sum_{m=1}^{\lfloor \frac{1}{k\mu} \rfloor} \delta_m u^m, u).$$

This transformation only shifts lines parallel to the z -axis, and has no effect on convexity in the z direction. Hence, denoting by γ^* the image of γ , we have

$$(3.7) \quad D_z^2 Q^*(z, u; \zeta_0) \leq -\epsilon'(|z|^{k-2} + |u|^{\frac{1}{\mu}(1-\frac{2}{k})})|\zeta_0|^2$$

on γ^* in a sufficiently small neighbourhood of the origin and $\epsilon' > 0$ sufficiently small. By homogeneity, it follows that F^* cannot be locally convex.

Now consider the second case, when f contains terms of weight strictly less than one. Then the leading weighted homogeneous term of F^* , which we denote $\hat{P}$, is harmonic in z . Let $a_{mol}z^m u^l$ be the nonzero monomial in this leading term with the lowest power of u , which dominates $D_z^2 \hat{P}$ along the z plane. Denote by $\nu = \frac{m}{k} + \mu l$ its weight, and let $k' = \nu k$. By harmonicity, there is a smooth curve γ in $\mathbb{C} \times \mathbb{R}$ and $\zeta_0 \neq 0$, such that in a neighbourhood of p

$$(3.8) \quad D_z^2 \hat{P}(z, u; \zeta_0) \leq -\epsilon'(|z|^{k'-2} + |u|^{\frac{1}{\nu}(1-\frac{2}{k'})})|\zeta_0|^2$$

for $(z, u) \in \gamma$ and a sufficiently small $\epsilon' > 0$. After applying S_0 , we use the same argument as in the first case. Consider terms of degree less or equal to $s + 2$ in F , where $s = (1 - \frac{2}{k'})\frac{1}{\nu}$, and denote them by T^* . Again, the same estimate holds for $D_z^2 T^*$ as for $D_z^2 \hat{P}$. After applying S_0 , we consider terms of weight less or equal to ν in F^{**} , denoted by Q^{**} . By (3.3) we obtain, modulo terms $o_{wt}(\nu)$

$$Q^{**}(z, \bar{z}, u) = T^*(z + \sum_{j=1}^{\lfloor \frac{1}{k\nu} \rfloor} \delta_j u^j, \bar{z} + \sum_{j=1}^{\lfloor \frac{1}{k\nu} \rfloor} \delta_j u^j, u).$$

On the image of γ , (3.8) holds for Q^{**} . By homogeneity, it follows that F^{**} cannot be locally convex. □

4. CONVEXIFIABILITY AND BASIC INVARIANTS

For applications of Propositions A and B we will need the explicit results obtained in [10]. For two even integers k, l and a positive real number a denote

$$M_a^{k,l} = \{(z, w) \in \mathbb{C}^2 \mid \operatorname{Im} w = P_a^{k,l}(z)\},$$

where

$$P_a^{k,l}(z, \bar{z}) = |z|^k + a|z|^{k-l} \operatorname{Re} z^l.$$

The Kohn-Nirenberg example is a hypersurface of this type, with $k = 8$, $l = 6$ and $a = \frac{15}{7}$.

Proposition C. $M_a^{k,l}$ is convex if and only if $a \leq \gamma_{lk}$, where

$$\gamma_{lk} = \frac{k}{l^2 - k}$$

if $l^2 \geq 3k - 2$ and

$$\gamma_{lk} = \sqrt{\frac{(4k - l^2 - 4)k^2}{(4k - 4)(k^2 - l^2)}}$$

if $l^2 \leq 3k - 2$. Moreover, if l is not a divisor of k , then this condition is equivalent to convexifiability of $M_a^{k,l}$.

Now we consider the general case, when the hypersurface is given by

$$(4.1) \quad v = P_1(z, \bar{z}) + o(|z|^k, u),$$

where $P_1(z, \bar{z})$ is a polynomial of the form (2.2). For such domains we define the basic invariants.

Definition 4.1. Let the model hypersurface to M at p be given by (2.3), and let l be an even integer, $0 < l < k$. We will call the real number $\kappa_l = \frac{|a_l|}{a_0}$ the Kohn-Nirenberg invariant of order l .

The numbers κ_l can be viewed as the 3-rd level local biholomorphic invariants, following the signature of the Levi form and the type of the point. The constant $\frac{15}{7}$ in the definition of D_0 is an invariant of order six.

In the general case we have a sufficient and a necessary condition for local convexifiability ([10], [11]).

Theorem 4.2. *Let the model at $p \in M$ be given by (2.3). If*

$$\sum_{j=2,4,\dots,k-2} \gamma_{jk}^{-1} \kappa_j < 1,$$

then M is locally convexifiable at p .

Theorem 4.3. *Let the model at $p \in M$ be given by (2.3). If M is locally convexifiable at p , then*

- (i) $\kappa_j \leq \gamma_{jk}$ for all $j > \frac{k}{2}$
- and
- (ii) $\kappa_j \leq 2\gamma_{jk}$ for all $j \leq \frac{k}{2}$.

5. EXAMPLES OF NONCONVEXIFIABLE DOMAINS WITH CONVEX MODELS

Using Propositions B and C we can find pseudoconvex domains which are not convexifiable, although their model domains at each point are convex.

As an example, consider a hypersurface M given by $v = F(z, \bar{z}, u)$, where

$$F(z, \bar{z}, u) = P_1(z, \bar{z}) + Q(z, \bar{z})u^s + o_{wt}(1),$$

and Q is a nonzero homogenous polynomial of degree $m < k$ without harmonic terms (both m and s are even). It is easy to verify that in this case $\mu = \frac{1}{s} - \frac{m}{ks}$ and $P(z, \bar{z}, u) = P_1(z, \bar{z}) + Q(z, \bar{z})u^s$ is homogeneous of weight one with respect to the weight vector $(\frac{1}{k}, \frac{1}{s} - \frac{m}{ks})$.

Since D_z^2 of $Q(z, \bar{z})u^s$ dominates D_z^2 of P along the u axis, we get from Proposition B the following condition. If M is convexifiable, then there exist $\alpha, \beta \in \mathbb{C}$ such that $P_1(z, \bar{z}) + \operatorname{Re} \alpha z^k$ and $Q(z, \bar{z}) + \operatorname{Re} \beta z^m$ are convex. Let us remark that for verifying this one can use the simpler complex form of the convexity condition - a function $f(z, \bar{z})$ is convex if and only if

$$|f_{z\bar{z}}| \leq f_{z\bar{z}}.$$

More explicitly, for $a > 0$ denote $P_a^{6,4}(z, \bar{z}) = |z|^6 + a|z|^2 \operatorname{Re} z^4$ and consider the hypersurface M_a defined by

$$v = |z|^8 + P_a^{6,4}(z, \bar{z})u^2 + |z|^2 u^8.$$

Using (3.3) we verify that $\mu = \frac{1}{k} = \frac{1}{8}$. The generalized model is given by $v = P(z, \bar{z}, u)$, where

$$P(z, \bar{z}, u) = |z|^8 + (|z|^6 + a|z|^2 \operatorname{Re} z^4)u^2.$$

To obtain the pseudoconvexity condition for M_a in terms of F , we take $r = F - v$ as the defining function and use the coordinate expression of the Levi form,

$$\mathcal{L} = |r_w|^2 r_{z\bar{z}} + |r_z|^2 r_{w\bar{w}} - 2\operatorname{Re}(r_{z\bar{w}} r_{\bar{z}w}),$$

which gives

$$\mathcal{L} = F_{z\bar{z}}(1 + |F_u|^2) + F_{uu}|F_z|^2 + 2\operatorname{Re} F_{zu}(F_u + i)F_{\bar{z}}.$$

Hence

$$\mathcal{L} = \frac{1}{4}\Delta F + o_{wt}(1 - \frac{2}{k}).$$

It follows that M_a is pseudoconvex if $\Delta P_a^{6,4} > 0$ away from the origin, which leads to $a < \frac{9}{5}$. In this case $l^2 = 3k - 2$ in Proposition C, which implies that M_a is not convexifiable if $a > \frac{3}{5}$. Hence for $\frac{3}{5} < a < \frac{9}{5}$ the hypersurface M_a is pseudoconvex and nonconvexifiable. On the other hand, the model at zero is the convex domain $v = |z|^8$, and all other points in a neighbourhood of zero are strongly pseudoconvex, hence the models are also convex.

REFERENCES

- [1] Catlin, D., *Boundary invariants of pseudoconvex domains*, Ann. Math. **120** (1984), 529–586.
- [2] D’Angelo, J., *Orders of contact, real hypersurfaces and applications*, Ann. Math. **115** (1982), 615–637.
- [3] S.S.Chern and J.Moser: *Real hypersurfaces in complex manifolds*, Acta Math. **133** (1974), p. 219–271
- [4] Diederich, K. and Fornæss, J. E., *Support functions for convex domains of finite type*, Math. Z. **230** (1999), 145–164.
- [5] Diederich, K. Fornæss, J. E. *Lineally convex domains of finite type: holomorphic support functions*, Manuscripta Math. **112** (2003), 403–431.
- [6] Diederich, K. and McNeal, J. D., *Pointwise nonisotropic support functions on convex domains*, Progress Math. **188** (2000), 184–192.
- [7] Fornæss, J. E. and Stenšones, B., *Lectures on Counterexamples in Several Complex Variables*, Princeton Univ. Press 1987.
- [8] Kohn, J. J., *Boundary behaviour of $\bar{\partial}$ on weakly pseudoconvex manifolds of dimension two*, J. Differential Geom. **6** (1972), 523–542.
- [9] Kohn, J. J. and Nirenberg, L., *A pseudoconvex domain not admitting a holomorphic support function*, Math. Ann. (1973), 265–268.
- [10] Kolář, M., *Convexifiability and supporting functions in $\mathbb{C}^2$* , Math. Res. Lett. **2** (1995), 505–513.
- [11] Kolář, M., *Necessary conditions for local convexifiability of pseudoconvex domains in $\mathbb{C}^2$* , Rend. Circ. Mat. Palermo **69** (2002), 109–116.
- [12] M.Kolář : *Normal forms for hypersurfaces of finite type in $\mathbb{C}^2$* , Math. Res. Lett. **12** (2005) p. 897–910
- [13] McNeal, J. D., *Estimates on the Bergman Kernels on Convex Domains*, Adv. Math. **109** (1994), 108–139.
- [14] McNeal, J. D., *Uniform subelliptic estimates on scaled convex domains of finite type*, Proc. Amer. Math. Soc. **130** (2002), 39–47 (electronic).

DEPARTMENT OF MATHEMATICAL ANALYSIS, MASARYK UNIVERSITY
JANÁČKOVO NÁM. 2A, 602 00 BRNO
E-mail: mkolar@math.muni.cz